Author Index

Adlard, M. W., see Orford, C. D. 245
Alawi, M., see Fayyad, M. 439
Arbutina, Ž. J., see Janjić, T. J. 465
Arco, A. Del, see Montes, M. 97
Avilés, F. X., see Burgos, F. J. 233
Bachman, W. J.
——and Stewart, J.T.
Optimization in photochemical reaction detection. Application to high-performance liquid chromatography–photolysis–electrochemical detection 121
Bächman, K.
——and Polzer, J.
Determination of tropospheric phosgene and other halocarbons by capillary gas chromatography 373
Balkom, C. A. A. van, see Geerdink, R. B. 275
Bennet, S., see Trumbore, C. N. 111
Berndt, H., see Kuropka, R. 380
Bernreuther, A.
——, Christoph, N. and Schreier, P.
Determination of enantiomeric composition of γ-lactones in complex natural matrices using multidimensional capillary gas chromatography 363
Brouwer, H.-J., see Geerdink, R. B. 275
Brown, D. W., see Krahn, M. M. 263
Burgos, F. J.
——, Pascual, R., Vendrell, J., Cuchillo, C. M. and Avilés, F. X.
The separation of pancreatic procarboxypeptidases by high-performance liquid chromatography and chromatofocusing 233
Burinsky, D. J., see Motto, M. G. 255
Burr, C. M., see Smith, R. M. 71, 85
Carr, J. W.
—— and Harris, J. M.
Temperature-induced changes in reversed-phase chromatographic surfaces. C_8 and C_9 polymeric ligands 135
Carté, B. K., see Poehland, B. L. 421
Casado, J., see Montes, M. 97
Ćelap, M. B., see Janjić, T. J. 465
——, see Tesic, Z. Lj. 471
Chase, H. A., see Livingston, A. G. 159
Christoph, N., see Bernreuther, A. 363
Cotter, M. L., see Motto, M. G. 255
Cuchillo, C. M., see Burgos, F. J. 233
Décout, J.-L.
——, Mouchel, B. and Lhomme, J.
Reversed-phase high-performance liquid chromatography, a tool for the study of bichromophoric systems including polymethylenic linking bridges 461
Del Arco, A., see Montes, M. 97
Do, L.
—— and Raulin, F.
Gas chromatography of Titan's atmosphere. I. Analysis of low-molecular-weight hydrocarbons and nitriles with a PoraPLOT Q porous polymer coated open-tubular capillary column 45
Dunphy, R., see Motto, M. G. 255
El-Ahmad, T., see Fayyad, M. 439
English, S., see Greenaway, W. 352
Eriksson, K.-O., see Hjertén, S. 175
Facchine, K. L., see Motto, M. G. 255
Fayyad, M.
——, Alawi, M. and El-Ahmad, T.
High-performance liquid chromatographic determination of phenoxyalkanoic acid herbicides using iron(II) 1,10-phenanthroline as mobile phase additive 439
Fournet, B., see Piot, J.-M. 221
Fujio, N., see Hatayama, T. 403
Funae, Y., see Hatayama, T. 403
Funazo, K.
——, Tanaka, M., Yasaka, Y., Takigawa, H. and Shono, T.
New ultraviolet labelling agents for high-performance liquid chromatographic determination of monocarboxylic acids 211
Geerdink, R. B.
——, Van Balkom, C. A. A. and Brouwer, H.-J.
Determination of phenoxyacid herbicides in water. Polymeric pre-column preconcentration and tetrabutylammonium ion-pair separation on a PRP-1 column 275
Gottstein, H. D.
——, Zook, M. N. and Kuć, J. A.
Detection and quantitation of oxalic acid by capillary gas chromatography 55
Gray, M. R., see Khorasheh, F. 1

Greenaway, W.
——, English, S., Wollenweber, E. and Whatley, F. R.
Series of novel flavanones identified by gas chromatography–mass spectrometry in bud exudate of *Populus fremontii* and *Populus maximowiczii* 352
Guillochon, D., see Piot, J.-M. 221
Günther, K., see Hünig, S. 387
Hamburg, P. F., see Motto, M. G. 255
Hardy, J. K., see Park, Y. J. 287
Harris, J. M., see Carr, J. W. 135
Hatayama, T.
——, Fujio, N., Yukioka, M., Funae, Y. and Kinoshita, H.
Separation of rat liver HSP70 and HSP71 by high-performance liquid chromatography with a hydroxylapatite column 403
Hayakawa, K.
——, Kitamoto, S., Okubo, N., Nakamura, S. and Miyazaki, M.
Determination of trace levels of total carbonate-carbon by indirect photometric ion chromatography with nitrogen purging 323
He, L.-Y.
——, Zhang, G.-D., Tong, Y.-Y., Sagara, K., Oshima, T. and Yoshida, T.
Reversed-phase ion-pair high-performance liquid chromatographic separation and determination of tropane alkaloids in Chinese solanaceous plants 428
Henry, R. J.
Rapid separation of α-amylases from barley by ion-exchange high-performance liquid chromatography on non-porous columns 397
Himberg, K.
Determination of anatoxin-a, the neurotoxin of *Anabaena flos-aquae* cyanobacterium, in algae and water by gas chromatography—mass spectrometry 358
Hirayama, N., see Maruo, M. 315
Hjertén, S.
——, Zelikman, I., Lindeberg, J. and Lederer, M.
High-performance adsorption chromatography of proteins on deformed non-porous agarose beads coated with insoluble metal compounds. II. Coating: aluminium and zirconium (hydr)oxide with stoichiometrically bound phosphate 187
——, Zelikman, I., Lindeberg, J. Liao, J.-l., Eriksson, K.-O. and Mohammad, J.
High-performance adsorption chromatography of proteins on deformed non-porous agarose beads coated with insoluble metal compounds. I. Coating: ferric oxyhydroxiide with stoichiometrically bound phosphate 175
Höcker, H., see Kuropka, R. 380
Houglum, J. E.
——, Larson, R. D. and Neal, R. M.
High-performance liquid chromatographic separation of indandione rodenticides 458
Hünig, S.
——, Klaunzer, N. and Günther, K.
Enantiomer separation of α-substituted γ-butyrolactones on the chiral polyacrylamide resin ChiraSpher® 387
Inoue, M., see Okada. T. 299
Ioneda, T.
Separation of homologues of methyl ester and 3-O-acetyl methyl ester derivatives of the corynomycolic acid fraction from *Corynebacterium pseudotuberculosis* 411
Izquierdo, C., see Montes, M. 97
Jackson, L. M., see Trumbore, C. N. 111
Janjić, T. J.
——, Milojković, D. M., Arbutina, Ž. J., Tešic, Ž. Lj. and Čelap, M. B.
Thin-layer chromatography on polyacrylonitrile. I. Effect of the *cis–trans* configuration of cobalt(III) complexes on their R_F values 465
——, see Tešić, Ž. Lj. 471
Jerzak, B., see Konopski, L. 477
Katano, M., see Matsunaga, H. 368
Kato, Y., see Yamasaki, Y. 391
Khorasheh, F.
——, Gray, M. R. and Selucky, M. L.
Correlation for Kováts retention index of C_9–C_{26} monoalkyl and polymethyl alkanes and alkenes 1
Kimata, K., see Ohtsu, Y. 147
Kinoshita, H., see Hatayama, T. 403
Kitamoto, S., see Hayakawa, K. 323
Kitamura, T., see Yamasaki, Y. 391
Klaunzer, N., see Hünig, S. 387
Kohama, A., see Nakagiri, I. 434
Komori, T.
—— and Morimoto, Y.
Isolation of the aromatic heptaenic antibiotics trichomycin A–F by high-performance liquid chromatography 416
Konopski, L.
—— and Jerzak, B.
2-Trichloromethylbenzimidazole, a selective chromogenic reagent for the detection of some azines on thin-layer plates (Letter to the Editor) 477
Koppenhoefer, B.
—— and Lin, B.
Thermodynamic properties of enantiomers of underivatized diols *versus* the cyclic carbonates in gas chromatography on Chirasil-Val 17
Koyama, J.-I., see Ohtsu, Y. 147

Krahn, M. M.
——, Wigren, C. A., Moore, L. K. and Brown, D. W.
High-performance liquid chromatographic method for isolating coprostanol from sediment extracts 263
Kuč, J. A., see Gottstein, H. D. 55
Kuhn, R.
—— and Wagner, H.
Application of free flow electrophoresis to the preparative purification of basic proteins from an *E. coli* cell extract 343
Kuroda, Y., see Nakagiri, I. 434
Kuropka, R.
——, Müller, B., Höcker, H. and Berndt, H.
Chiral stationary phases via hydrosilylation reaction of N-acryloylamino acids. I. Stationary phase with one chiral centre for high-performance liquid chromatography and development of a new derivatization pattern for amino acid enantiomers 380
Kuwamoto, T., see Maruo, M. 315
Laan, A. van der, see Vertommen, M. H. 452
Larson, R. D., see Houglum, J. E. 458
Lederer, M., see Hjertén, S. 187
Lhomme, J., see Décout, J.-L. 461
Liao, J.-l., see Hjertén, S. 175
Lin, B., see Koppenhoefer, B. 17
Lindeberg, J., see Hjertén, S. 175, 187
Livingston, A. G.
—— and Chase, H. A.
Preparation and characterization of adsorbents for use in high-performance liquid affinity chromatography 159
McGarvey, B. D.
Liquid chromatographic determination of N-methylcarbamate pesticides using a single-stage post-column derivatization reaction and fluorescence detection 445
Malinar, M. J., see Tešić, Ž. Lj. 471
Mallet, C.
—— and Mallet, V. N.
Conversion of a conventional packed-column gas chromatograph to accommodate megabore columns. I. Evaluation of the system for organophosphorus pesticides 27
—— and Mallet, V. N.
Conversion of a conventional packed-column gas chromatograph to accommodate megabore columns. II. Determination of organophosphorus pesticides in environmental water 37
Mallet, V. N., see Mallet, C. 27, 37
Maruo, M.
——, Hirayama, N. and Kuwamoto, T.
Ion chromatographic elution behaviour and prediction of the retention of inorganic monovalent anions using a phosphate eluent 315
Matsunaga, H.
——, Katano, M., Yamamoto, H., Mori, M., Takata, K. and Nishi, M.
Determination of panaxytriol, a new type of tumour growth inhibitor from Panax ginseng, by capillary gas chromatography 368
Milojković, D. M., see Janjić, T. J. 465
Miyakoshi, M., see Okada, T. 299
Miyazaki, M., see Hayakawa, K. 323
Mohammad, J., see Hjertén, S. 175
Mohanty, M., see Patnaik, L. N. 331
Montes, M.
——, Usero, J. L., Del Arco, A., Izquierdo, C. and Casado, J.
Free energy correlations: dead volume and the reversed-phase high-performance liquid chromatographic capacity factor in the interaction index model. A discussion and application to a nitrosamine series 97
Moore, L. K., see Krahn, M. M. 263
Mori, M., see Matsunaga, H. 368
Morimoto, Y., see Komori, T. 416
Motto, M. G.
——, Facchine, K. L., Hamburg, P. F., Burinsky, D. J., Dunphy, R., Oyler, A. R. and Cotter, M. L.
Separation and identification of retinoic acid photoisomers 255
Mouchel, B., see Décout, J.-L. 461
Müller, B., see Kuropka, R. 380
Nakabayashi, Y., see Okada, T. 299
Nakagiri, I.
——, Suzuki, K., Shiaku, Y., Kuroda, Y., Takasu, N. and Kohama, A.
Rapid quantification of paraquat and diquat in serum and urine using high-performance liquid chromatography with automated sample pretreatment 434
Nakamura, K., see Ohtsu, Y. 147
Nakamura, S., see Hayakawa, K. 323
Nakata, O., see Ohtsu, Y. 147
Nakatani, S., see Yamasaki, Y. 391
Neal, R. M., see Houglum, J. E. 458
Nishi, M., see Matsunaga, H. 368
Ohtsu, Y.
——, Shiojima, Y., Okumura, T., Koyama, J.-I., Nakamura, K., Nakata, O., Kimata, K. and Tanaka, N.
Performance of polymer-coated silica C_{18} packing materials prepared from high-purity silica gel. The suppression of undesirable secondary retention processes 147

Okada, T.
——, Sugata, K., Nakabayashi, Y., Teraoka, K., Miyakoshi, M. and Inoue, M.
Exchange of the cation and anion of the sample (sodium or potassium chloride) with the cation and anion of the eluent (sodium or potassium phosphate buffer) and their elution, from a Sephadex G-15 column, in separate fractions 299
Okubo, N., see Hayakawa, K. 323
Okumura, T., see Ohtsu, Y. 147
Orford, C. D.
——, Perry, D. and Adlard, M. W.
High-performance liquid chromatographic determination of δ-(L-α-aminoadipyl)-L-cysteinyl-D-valine in complex media by precolumn derivatisation with dansylaziridine 245
Oshima, T., see He, L.-Y. 428
Oyler, A. R., see Motto, M. G. 255
Park, Y. J.
—— and Hardy, J. K.
Reversed-phase high-performance liquid chromatography of metal–benzylpropionitrile dithiocarbamate complexes 287
Pascual, R., see Burgos, F. J. 233
Patnaik, L. N.
——, Pattanaik, B. N., Mohanty, M. and Satapathy, A.
Thin-layer chromatographic behaviour of some styryl cyanine dyes derived from pyridine 331
Pattanaik, B. N., see Patnaik, L. N. 331
Perry, D., see Orford, C. D. 245
Piot, J.-M.
——, Guillochon, D., Zhao, Q., Ricart, G., Fournet, B. and Thomas, D.
Identification of peptides, from a peptic haemoglobin hydrolysate produced at pilot-plant scale, by high-performance liquid chromatography and mass spectrometry 221
Poehland, B. L.
——, Troupe, N., Carté, B. K. and Westley, J. W.
Reversed-phase high-performance liquid chromatographic assay for camptothecin and related alkaloids 421
Polzer, J., see Bächmann, K. 373
Raulin, F., see Do, L. 45
Ricart, G., see Piot, J.-M. 221
Sagara, K., see He, L.-Y. 428
Sanagi, M. M., see Smith, R. M. 63
Satapathy, A., see Patnaik, L. N. 331
Sawai, H.
Preparation of several types of RPC-5-like resins and their use for the separation of oligonucleotides and mononucleotides by high-performance liquid chromatography 201
Schreier, P., see Bernreuther, A. 363
Selucky, M. L., see Khorasheh, F. 1
Shiaku, Y., see Nakagiri, I. 434
Shiojima, Y., see Ohtsu, Y. 147
Shono, T., see Funazo, K. 211
Smith, R. M.
—— and Burr, C. M.
Retention prediction of analytes in reversed-phase high-performance liquid chromatography based on molecular structure. III. Monosubstituted aliphatic compounds 71
—— and Burr, C. M.
Retention prediction of analytes in reversed-phase high-performance liquid chromatography based on molecular structure. IV. Branched and unsaturated alkylbenzenes 85
—— and Sanagi, M. M. Supercritical fluid chromatography of barbiturates 63
Stewart, J. T., see Bachman, W. J. 121
Sugata, K., see Okada, T. 299
Suzuki, K., see Nakagiri, I. 434
Takasu, N., see Nakagiri, I. 434
Takata, K., see Matsunaga, H. 368
Takigawa, H., see Funazo, K. 211
Tanaka, M., see Funazo, K. 211
Tanaka, N., see Ohtsu, Y. 147
Teraoka, K., see Okada, T. 299
Tešić, Ž. Lj.
——, Janjić, T. J., Malinar, M. J. and Ćelap, M. B.
Effect of the chelate ring size of diamine cobalt (III) complexes on their R_F values obtained by paper chromatography 471
——, see Janjić, T. J. 465
Thomas, D., see Piot, J.-M. 221
Thompson, A., see Trumbore, C. N. 111
Tong, Y.-Y., see He, L.-Y. 428
Troupe, N., see Poehland, B. L. 421
Trumbore, C. N.
——, Jackson, L. M., Bennett, S. and Thompson, A.
High-speed analytical sensor for in-line monitoring of dissolved analytes flowing in a tube empoying a combination of limited diffusion, laminar flow and plug solvent injections 111
Usero, J. L., see Montes, M. 97
Van Balkom, C. A. A., see Geerdink, R. B. 275
Van der Laan, A., see Vertommen, M. H. 452
Veenendaal-Hesselman, H. M., see Vertommen, M. H. 452
Vendrell, J., see Burgos, F. J. 233
Vertommen, M. H.
——, Van der Laan, A. and Veenendaal-Hesselman, H. M.
High-performance liquid chromatographic screening method for low levels of nicarbazin in eggs with off-line cartridge sample clean-up 452

Wagner, H., see Kuhn, R. 343
Westley, J. W., see Poehland, B. L. 421
Whatley, F. R., see Greenaway, W. 352
Wigren, C. A., see Krahn, M. M. 263
Wollenweber, E., see Greenaway, W. 352
Yamamoto, H., see Matsunaga, H. 368
Yamasaki, Y.
——, Kitamura, T., Nakatani, S. and Kato, Y. Recovery of proteins and peptides with nanogram loads on non-porous packings 391
Yasaka, Y., see Funazo, K. 211
Yoshida, T., see He, L.-Y. 428
Yukioka, M., see Hatayama, T. 403
Zelikman, I., see Hjertén, S. 175
Zelikman, I., see Hjertén, S. 187
Zhang, G.-D., see He, L.-Y. 428
Zhao, Q., see Piot, J.-M. 221
Zook, M. N., see Gottstein, H. D. 55